*Fluid Phase Equilibria*, 58 (1990) 337–338
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands



## Contents Volume 58 (1990)

*Volume 58, Nos. 1–2*

Polar pressure of water and simplified augmented BACK equation of state
M.J. Lee and K.C. Chao (West Lafayette, IN, U.S.A.) . . . . . 1– 12

Evaluation of the simplified perturbed hard chain theory (SPHCT) for prediction of phase behavior of n-paraffins and mixtures of n-paraffins with ethane
K.A.M. Gasem and R.L. Robinson, Jr. (Stillwater, OK, U.S.A.) . . . . . 13– 33

Prediction of the thermophysical properties of fluid methane from the SSR–MPA potential
M. Luckas, M. Ripke and K. Lucas (Duisburg, F.R.G.) . . . . . 35– 50

A renormalization group background transform lattice model for binary liquid–liquid equilibria
P.D. McMahon (Madison, WI, U.S.A.) . . . . . 51– 64

The critical lines of the van der Waals equation for binary mixtures around the van Laar point
P.H.E. Meijer, I.L. Pegg (Washington, DC, U.S.A.), J. Aronson (Gaithersburg, MD, U.S.A.) and M. Keskin (Kayseri, Turkey) . . . . . 65– 80

Local composition in ternary Lennard-Jones liquid mixtures: molecular dynamics simulation and a simple correlation
J.A. Abusleme, K. Nakanishi (Kyoto, Japan) and J.H. Vera (Montreal, PQ, Canada) . . . . . 81– 92

A new mixing rule for the Patel–Teja equation of state. Study of vapor–liquid equilibria
H. Huang (Beijing, China) . . . . . 93–115

Estimation of multiple binary interaction parameters in equations of state using VLE data. Application to the Trebble–Bishnoi equation of state
P. Englezos, N. Kalogerakis, M.A. Trebble and P.R. Bishnoi (Calgary, Alb., Canada) . . . . . 117–132

Extended phase rule for non-reactive, multiphase, multicomponent chemical systems
L.J. Van Poolen (Boulder, CO, U.S.A.) . . . . . 133–145

Measurement and correlation of liquid–liquid equilibria of methanol–alkane–aromatic compound ternary systems
H. Higashiuchi, Y. Sakuragi (Kurume, Japan), Y. Arai and M. Nagatani (Fukuoka, Japan) . . . . . 147–157

Solubility of 15 non-polar gases (He, Ne, Ar, Kr, Xe, $H_2$, $D_2$, $N_2$, $O_2$, $CH_4$, $C_2H_4$, $C_2H_6$, $CF_4$, $SF_6$ and $CO_2$) in cycloheptanone
M.A. Gallardo, M.C. López, J.S. Urieta and C. Gutiérrez Losa (Zaragoza, Spain) . . . . . 159–172

Excess molar volumes of binary mixtures of benzene + n-propylbenzene, + n-butylbenzene and + n-hexylbenzene at 298.15, 308.15 and 318.15 K
D.V.S. Jain and N.S. Dhar (Chandigarh, India) . . . . . 173–180

Elsevier Science Publishers B.V.

Determination of vapor–liquid equilibria by total pressure measurements for four binary silicon compound systems: tetramethoxy silane + *p*-xylene, tetramethoxy silane + *n*-nonane, tetraethoxy silane + *p*-xylene, and tetraethoxy silane + *n*-nonane
M. Kato, H. Tanaka and K. Aizawa (Fukushima, Japan) . . . . . 181–189

Total vapour pressure measurements and excess Gibbs energies for the binary systems methanol + ethanol, ethanol + 2-propanol, benzene + cyclohexane, benzene + carbon tetrachloride and benzene + ethanol at 303.15 and 313.15 K
J. Zielkiewicz (Gdansk, Poland), P. Oracz and S. Warycha (Warsaw, Poland) . 191–209

Solubility of normal alkanoic acids in selected organic binary solvent mixtures. Negative synergetic effect
U. Domańska and K. Kniaź (Warsaw, Poland) . . . . . 211–227

*Letter to the Editor*

On composition-dependent interaction coefficients
M.L. Michelsen (Lyngby, Denmark) and H. Kistenmacher (Höllriegelskreuth, F.R.G.) . . . . . 229–230

*Volume 58, No. 3*

A new generalized equation for predicting volumes of compressed liquids
C.-H. Chang and X.-M. Zhao (Tianjin, China) . . . . . 231–238

The significant structure model equation of state extended to mixtures
K. Suzuki, H. Sue (Chiba, Japan), H. Inomata, K. Arai and S. Saito (Sendai, Japan) . . . . . 239–264

The simplified perturbed anisotropic chain theory: a new equation of state for polar fluids
G. Ciocca and I. Nagata (Kanazawa, Japan) . . . . . 265–282

The molecular statistical theory of infinitely dilute solutions based on the ion–dipole model with Lennard-Jones interaction
S.N. Lvov, V.A. Umniashkin, A. Sharygin (Leningrad, U.S.S.R.) and M.F. Holovko (Lvov, U.S.S.R.) . . . . . 283–305

Activity coefficient data and their correlation for tributyl phosphate–hydrocarbon and uranyl nitrate tributyl phosphate complex–hydrocarbon solutions
J.-D. Li, Y.-G. Li, J. Chen, J-F. Lu and T. Teng (Beijing, China) . . . . . 307–318

Excess volumes for binary liquid mixtures of 1,4-dioxane with methylene chloride, 1,2-dichloroethane, trichloroethylene, tetrachloroethylene and cyclohexane at various temperatures
J. Nath and Rashmi (Gorakhpur, India) . . . . . 319–324

Mutual solubility of binary *trans*-decalin + , and n-decane + polar component mixtures
M. Antosik, A. Stafiej and R. Stryjek (Warsaw, Poland) . . . . . 325–333

*Author Index* . . . . . 335

*Contents of Volume 58 (1990)* . . . . . 337–338